

¹The CDC defines immune-compromised as individuals who have:


- Been receiving active cancer treatment for tumors or cancers of the blood
- Received an organ transplant and are taking medicine to suppress the immune system
- Received a stem cell transplant within the last 2 years or are taking medicine to suppress the immune system
- Moderate or severe primary immunodeficiency (such as DiGeorge syndrome, Wiskott-Aldrich syndrome)
- Advanced or untreated HIV infection
- Active treatment with high-dose corticosteroids or other drugs that may suppress your immune response

² Primary series dose volumes are as follows:

- Pediatric Pfizer (ages 5-11) 0.2mL
- Pfizer (ages 12+) 0.3mL
- Moderna 0.5mL
- Janssen 0.5mL

³ It is recommended that all individuals should receive a booster dose of an mRNA COVID-19 vaccine (Pfizer or Moderna) two months after their Janssen (Johnson & Johnson) vaccine.

⁴ All individuals are recommended to receive a primary vaccination series and booster dose of an mRNA COVID-19 vaccine (Pfizer or Moderna) over the use of the Janssen (Johnson & Johnson) COVID-19 vaccine. This change in guidance comes after reviewing data finding an increased risk of Thrombosis with Thrombocytopenia associated with those who have received the Janssen COVID-19 vaccine.



¹All individuals are recommended to receive a primary vaccination series and booster dose of an mRNA COVID-19 vaccine (Pfizer or Moderna) over the use of the Janssen COVID-19 vaccine. This change in guidance comes after reviewing data finding an increased risk of Thrombosis with Thrombocytopenia associated with those who have received the Janssen COVID-19 vaccine.

²Booster dose volumes are as follows:

- Pediatric Pfizer (ages 5-11) 0.2mL
- Pfizer (ages 12+) 0.3mL
- Moderna 0.5mL
- Janssen 0.5mL

For additional questions, please refer to the [CDC guidelines](#) on vaccine booster eligibility.