

Update on Monoclonal Antibody Shipment

The distribution of allocated Regeneron moving forward may come from different concentrations and packaging every week. Please be diligent in noting correct dosage when mixing infusions and understand that some packaging contains more than one course of the medication.

Each REGEN-COV dose pack contains a sufficient number of vials of casirivimab [REGN10933] and imdevimab [REGN10987] to prepare up to two treatment doses (600 mg of casirivimab and 600 mg of imdevimab). Refer to Table listed below.

Dose Pack Size	Dose Pack Components	Concentration
2 Cartons	1 casirivimab REGN10933 (NDC 61755-024-01)	1,332 mg/11.1 mL (120 mg/mL)
	1 imdevimab REGN10987 (NDC 61755-025-01)	1,332 mg/11.1 mL (120 mg/mL)
8 Cartons	4 casirivimab REGN10933 (NDC 61755-026-01)	300 mg/2.5 mL (120 mg/mL)
	4 imdevimab REGN10987 (NDC 61755-027-01)	300 mg/2.5 mL (120 mg/mL)
5 Cartons	1 casirivimab REGN10933 (NDC 61755-024-01)	1,332 mg/11.1 mL (120 mg/mL)
	4 imdevimab REGN10987 (NDC 61755-027-01)	300 mg/2.5 mL (120 mg/mL)

5 Cartons	4 casirivimab REGN10933 (NDC 61755-026-01)	300 mg/2.5 mL (120 mg/mL)
	1 imdevimab REGN10987 (NDC 61755-025-01)	1,332 mg/11.1 mL (120 mg/mL)

It is expected that facilities are conscious of the number of doses they have and understand that some packaging will result in two courses of Regeneron once reconstituted. When scheduling Regeneron treatment facilities are should be cognizant of treatment appointment times to avoid wastage due to expiration of the treatment after reconstitution.

If immediate administration is not possible, store the diluted casirivimab and imdevimab infusion solution in the refrigerator between 2°C to 8°C (36°F to 46°F) for no more than 36 hours or at room temperature up to 25°C (77°F) for no more than 4 hours. If refrigerated, allow the infusion solution to equilibrate to room temperature for approximately 30 minutes prior to administration.

When doses are reconstituted, and the second course will not be utilized, the unused course may be sent to another facility for use. It is expected that facilities will reach out to surrounding health care facilities to see if they will have a need for the reconstituted dose to avoid wastage.

If assistance will be needed to transport reconstituted vaccine, please call 701-328-0707, we are available 24/7 to assist with distribution as needed.

#

Categories of Health Alert Network messages:

Health Alert Requires immediate action or attention; highest level of importance

Health Advisory May not require immediate action; provides important information for a specific incident or situation

Health Update Unlikely to require immediate action; provides updated information regarding an incident or situation

HAN Info Service Does not require immediate action; provides general public health information

##This message was distributed to state and local health officers, state and local epidemiologists, state and local laboratory directors, public information officers, HAN coordinators, and clinician organizations##