


# Measles Vaccine Recommendations

May 20, 2019



Note: There is no recommendation for an adult catch-up or vaccination campaign among adults or individuals in non-affected areas to prevent measles.

**Contraindications to receiving a dose of MMR:**

- Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to vaccine component
- Have a weakened immune system due to disease (such as cancer or HIV/AIDS) or medical treatments (such as radiation, immunotherapy, steroids or chemotherapy)
- Pregnancy
- Have a parent, brother, or sister with a history of immune system problems

**†High risk groups recommended to receive 2 doses of MMR (or have presumptive evidence of immunity\*)**

- International travelers
  - Adults and children ages 12 months or older: 2 doses of MMR
  - Children ages 6 to 11 months: 1 dose of MMR
- Healthcare personnel (even if born before 1957)
- Student at post-high school educational institutions (North Dakota colleges and universities require two doses of MMR)

**\*Presumptive evidence of immunity includes any of the following (documented, rather than self-reported):**

- Written documentation of age appropriate vaccination
- Laboratory confirmation of past infection or had blood tests that show you are immune to measles
- Born before 1957

‡ For complete recommendations visit: [www.cdc.gov/vaccines/schedules/index.html](http://www.cdc.gov/vaccines/schedules/index.html)